

Three reflections

Waldemar Pompe
University of Warsaw, Poland
pompe@mimuw.edu.pl

Abstract

We present several problems that can be solved in a very short way using properties of a glide reflection. In our configurations the glide reflection will be obtained as a composition of three reflections.

I dedicate this paper to the memory of Professor Edmund Puczyłowski.

1. Preliminary results

Consider a reflection about line x in a plane. For simplicity, we shall use for this reflection the same notation “ x ”; it should be always clear from the context, whether “ x ” means “line x ” or “the reflection about line x ”. Similarly, if v is a vector, we will also denote by v the translation by vector v .

In this paper we are going to compose reflections and other isometries. As usual, by a composition “ gf ” of two mappings f and g we mean the mapping defined by $(gf)(X) = (g \circ f)(X) = g(f(X))$.

The following theorem is known in the theory of geometric transformations (see [1, Theorem 3.31]).

Theorem 1.1

Let a, b, c be three lines in a plane (see Fig. 1). Then there exist a unique line d and a unique vector v parallel to d , such that

$$cba = dv.$$

Moreover, lines a, b, c are concurrent if and only if $v = 0$.




Fig. 1

Of course, general isometries f and g don't commute, i.e. $fg \neq gf$, but if vector v is parallel to line d , then it is immediately clear that they do: $dv = vd$.

The mapping $f = dv = vd$ with $d \parallel v$ is called a *glide reflection*, line d — the *mirror* or *axis* of f and v — the vector of f .

It follows from Theorem 1.1 that if lines a, b, c are concurrent, then cba reduces to a single reflection d . It turns out line d can be described in a simple geometric way. Namely, if lines a, b , and c are not parallel (see Fig. 2), then line d is determined by

$$(1) \quad \sphericalangle(d, c) = \sphericalangle(a, b).$$

Here $\sphericalangle(x, y)$ denotes the oriented angle between lines x and y , which is defined as an angle of rotation taking line x onto a line parallel to y . It is easy to see that such an angle is defined up to 180° (the same angles differ by an integer multiple of 180°).

To see that (1) implies $d = cba$ denote by α the angles given in equality (1). Then both cd and ba are rotations with the same center $O = a \cap b$ and the same angle 2α , so they are equal mappings. From this equality we immediately obtain $d = cba$, as cc is the identity mapping.

Line d satisfying equality (1) is called an *isogonal line* to b in the angle formed by lines a and c . Note that line d can be also obtained from b by reflecting it about one of the bisectors of the angles determined by lines a and c .




Fig. 2




Fig. 3

Similarly, we may describe line d , if lines a, b, c are parallel (see Fig. 3). Namely, if w is a vector perpendicular to a and b that moves line a to line b , then line d should be chosen in a way that w moves line d to line c . Then both cd and ba are translations by vector $2w$, implying $cd = ba$, that is $d = cba$.

Let's get back to Theorem 1.1. There is also a geometric way to describe line d and vector v , if lines a, b, c are not concurrent and form a triangle. For this purpose we will need to introduce a *signed perimeter of an orthic triangle*.

Definition 1.2

Let ABC be a triangle. Denote by D, E, F the feet of the altitudes of triangle ABC dropped from vertices A, B, C , respectively (see Fig. 4 and Fig. 5). A *signed perimeter of the orthic triangle* of triangle ABC , denoted by $\sigma(ABC)$, is defined by:

$$\sigma(ABC) = \begin{cases} DE + EF + FD, & \text{if } ABC \text{ is acute-angled,} \\ -DE + EF + FD, & \text{if } \angle ACB \geq 90^\circ. \end{cases}$$




Fig. 4




Fig. 5

Theorem 1.3

Let a, b, c be lines that determine triangle ABC with $A = b \cap c, B = c \cap a, C = a \cap b$ and $\angle ABC \neq 90^\circ$ (see Fig. 6). Denote by D, E, F the feet of the altitudes of triangle ABC taken from vertices A, B, C , respectively. If line d and vector v parallel to d are determined by the condition

$$cba = dv,$$

then d coincides with line DF and $|v| = \sigma(ABC)$.




Fig. 6

Theorem 1.3 is known and can be found in the literature, e.g. [3, Section 19, Problem 13]. A similar theorem for spherical triangles can be found in [2].

Proof

The proof uses the following well-known fact from the triangle geometry: The altitudes of a triangle are the angle bisectors of the angles of the orthic triangle. More precisely, if ABC is an acute-angled triangle with AD, BE, CF as altitudes, then $\angle FEB = \angle DEB$ (see Fig. 7). The proof follows immediately from the observation that $ABDE$ and $BCEF$ are cyclic quadrilaterals. Similar formulas hold, if triangle ABC is obtuse-angled.




Fig. 7




Fig. 8

We will also use the following simple observation about glide reflections. If a glide reflection $f = vd$, where d is the axis, and v is the vector of f , maps point X to X' , then the midpoint Y of segment XX' lies on axis d (see Fig. 8). Moreover, if Z is the foot of the perpendicular from X onto line d , then

$$\overrightarrow{ZY} = \frac{1}{2}v.$$

We turn to the proof of Theorem 1.3. For simplicity, we assume that triangle ABC is acute-angled (see Fig. 9). The proof for obtuse-angled triangles is almost the same, though diagrams are a bit different (see Fig. 10, 11).




Fig. 9

Set $f = cba$. From Theorem 1.1 we know that $f = dv$, where d is a line and v is a vector parallel to d .

Let K and L be the reflections of points C and A in lines c and a , respectively. Since $f = cba$, we infer that $f(C) = K$ and $f(L) = A$. Thus D and F are the midpoints of segments $Cf(C)$ and $Lf(L)$, so D and F belong to line d . This means that line d coincides with line DF .




Fig. 10




Fig. 11

Denote by P and Q the reflections of point E in lines a and c , respectively. Using the fact that we have mentioned at the beginning of the proof, we conclude that points P and Q lie on DF . Moreover f maps point P to Q , and since P lies on d , it implies $v = \overrightarrow{PQ}$. Therefore,

$$|v| = |\overrightarrow{PQ}| = \sigma(ABC),$$

which completes the proof.

Case $a \perp c$ is covered by the next theorem.

Theorem 1.4

Let a, b, c be lines that determine triangle ABC with $A = b \cap c, B = c \cap a, C = a \cap b$ and $\angle ABC = 90^\circ$ (see Fig. 12). Denote by E the foot of the altitude of triangle ABC taken from vertex B . If line d and vector v parallel to d are determined by the condition

$$cba = dv,$$

then d coincides with the tangent line to the circumcircle of triangle ABC at point B . Moreover, $|v| = \sigma(ABC) = 2BE$.

Proof

Denote by d the tangent line to the circumcircle of triangle ABC at point B . Let M be the midpoint of segment AC and let P be the foot of the perpendicular from C onto d . Moreover, denote by x and y lines PC and BM , respectively. Then x and y are parallel, since they are perpendicular to d .




Fig. 12

Moreover, $\sphericalangle(c, d) = \sphericalangle(b, a) = \sphericalangle(a, y) = \sphericalangle(a, x)$. It follows that $ax = ba$, so $x = aba$. Therefore, $yx = (ya)ba = dcba$, so $d(yx) = cba$. Setting

$$v = 2\overrightarrow{PB},$$

we obtain $dv = cba$.

Finally, $|v| = 2PB = 2BE = \sigma(ABC)$, which completes the proof.

We conclude the preliminary results with the following observation about glide reflections.

Remark 1.5

Let d be a line and let v a vector parallel to d (see Fig. 13). Moreover, let b be a line. Then $b(dv)b$ is a glide reflection, whose axis d' and vector v' are obtained from d and v , respectively, by reflection about line b .




Fig. 13

For a proof simply observe that the mappings $b(dv)b$ and $d'v'$ act on a sample point X in the same way.

2. Applications

Problem 2.1

Let $ABCD$ be an arbitrary quadrilateral. The perpendicular bisectors of segments AB, BC, CD bound triangle PQR , as shown in Figure 14. Points K and L are the feet of the altitudes of triangle PQR taken from vertices Q and R , respectively. Let M be the midpoint of side AD . Prove that points K, L , and M are collinear.




Fig. 14

Solution

Denote by a, b, c the perpendicular bisectors of segments AB, BC, CD , respectively. Then $cba = dv$, where line d coincides with KL and v is a vector parallel to KL . Observe now that cba maps A to D , so the midpoint M of AD lies on d . This completes the proof.

Problem 2.2

Let $ABCD$ be a convex quadrilateral (see Fig. 15). The bisectors x, y, z of angles $\angle A, \angle B, \angle C$, respectively, bound triangle T . Let X and Z be the feet of the altitudes of T dropped onto lines x and z , respectively. Line XZ meets side AD at point P . Prove that:

- (a) $AP + BC = AB + CD + DP$;
- (b) $\sigma(T) = 2DP \cdot \cos \alpha$, where $\alpha = \angle XPA$;

Solution

Reflecting line AD about lines x, y , and z , sequentially, we obtain lines AB, BC , and CD . So if we reflect point P about lines x, y, z , sequentially, we get points Q, R, S with Q on AB, R on BC , and S on CD (see Fig. 15). Moreover, $AP = AQ, BQ = BR$, and $CR = CS$.

Set $f = zyx$. By Theorem 1.3, we know that $f = dv$, where d coincides with line XZ and v is a vector parallel to d with $|v| = \sigma(T)$. Therefore, since P lies on d , point $S = f(P)$ must also belong to d and $\overrightarrow{PS} = v$.

Also, f maps line AD onto CD . It means line CD is obtained from AD by a reflection in line d followed by a translation. Therefore line d makes equal angles α with lines AD and CD . Thus we obtain

$$\sigma(T) = |v| = PS = 2PD \cos \alpha$$

which is (b).




Fig. 15

To prove (a), observe that

$$\begin{aligned} AP + BC &= AP + CR - BR = AQ + CS - BQ = AB + CS \\ &= AB + CD + DP. \end{aligned}$$

This completes the proof.

The next example is a very well-known theorem about the existence of the isogonal point in a triangle. The proof we are going to present is also known (see [3, Theorem 20.12] or [5]).

Problem 2.3

Let ABC be any triangle and let P be any point (see Fig. 16). Let x' , y' , z' be isogonal lines to lines AP , BP , CP in the angles $\angle A$, $\angle B$, $\angle C$, respectively. Prove that lines x' , y' , and z' are concurrent.

Solution

Denote by a , b , c lines BC , CA , AB , respectively, and by x , y , z lines AP , BP , CP , respectively.

Lines x , y , z are concurrent, so by Theorem 1.1 the mapping zyx is a reflection d . We want to prove that $z'y'x'$ is also a reflection. But

$$x' = cxb, \quad y' = ayc, \quad z' = bza.$$

Therefore, we obtain

$$z'y'x' = (bza)(ayc)(cxb) = b(zyx)b = bdb = d',$$

where d' is a line obtained from d in reflection about line b (see Remark 1.5). Thus $z'y'x'$ is a reflection, meaning that x' , y' , and z' concur. This completes the proof.




Fig. 16

From the above proof it follows that a line isogonal to y in angle $\angle(x, z)$ (which is d) and a line isogonal to y' in angle $\angle(x', z')$ (which is d') are symmetric to each other with respect to line b (see Fig. 17).

The next problem was proposed in 1952 by Victor Thébault for the American Mathematical Monthly (Problem 4470). The Proposer's published solution was based on trigonometric formulas [4].




Fig. 17

Problem 2.4

Let ABC be a triangle. Assume lines x, y, z passing through vertices A, B, C , respectively, bound triangle T . Lines x', y', z' are isogonal to lines x, y, z in angles A, B, C , respectively, of the triangle ABC . Assume that lines x', y', z' bound triangle T' . Prove that the signed perimeters of the orthic triangles of T and T' are equal, i.e. $\sigma(T) = \sigma(T')$.




Fig. 18

Solution

Consider the mapping $f = zyx$. From Theorem 1.1 we know that mapping f can be reduced to dv , where d is a line and v is a vector parallel to d .

Denote by a, b, c lines BC, CA, AB , respectively. Since x', y' , and z' are lines isogonal to x, y , and z in the corresponding angles of triangle ABC , we have:

$$x' = cxb, \quad y' = ayc, \quad z' = bza.$$

Therefore, we get

$$z'y'x' = (bza)(ayc)(cxb) = b(zyx)b = b(dv)b = d'v',$$

where d' is a line and v' a vector obtained from d and v , respectively, by reflection about line b (see Remark 1.5). Therefore, from Theorem 1.3 we immediately get

$$\sigma(T') = |v'| = |v| = \sigma(T),$$

which completes the proof.




Fig. 19

Since lines d and d' are symmetric to each other with respect to line b , we have also solved the following problem.

Problem 2.5

Given triangle ABC , construct triangles T and T' as in Problem 2.4 Let D and F be the feet of the altitudes of triangle T taken onto lines x and z , respectively (see Fig. 19). Similarly, D' and F' are the feet of the altitudes of triangle T' taken onto lines x' and z' , respectively. Prove that lines DF and $D'F'$ are symmetric to each other with respect to line AC . In particular, lines DF and $D'F'$ meet at point lying on line AC .

Problem 2.6.

Let $ABCDEF$ be a convex hexagon with $\angle B + \angle D + \angle F = 360^\circ$ (see Fig. 20). Denote by x, y, z, x', y', z' the perpendicular bisectors of segments AB, BC, CD, DE, EF, FA , respectively. Lines x, y , and z bound triangle T , while lines x', y' , and z' bound triangle T' . Finally, denote by K, L the feet of the altitudes of T taken onto lines x, z , respectively. Similarly, K', L' are the feet of the altitudes of T' taken onto lines x', z' , respectively. Prove that:

- (a) points K, L, K', L' are collinear;
- (b) $\sigma(T) = \sigma(T')$.




Fig. 20

Solution

Consider the mapping $f = z'y'x'zyx$. Observe that $yx, x'z, z'y'$ are rotations about angles

$$2\angle(x, y) = 360^\circ - 2\angle B, \quad 2\angle(z, x') = 360^\circ - 2\angle D, \quad 2\angle(y', z') = 360^\circ - 2\angle F,$$

respectively. Therefore, since $\angle B + \angle D + \angle F = 360^\circ$, we infer that the above angles sum up to 360° , so mapping $f = (z'y')(x'z)(yx)$ is a translation. But since A is a fixed point of f , it follows that f is an identity.

Therefore, we obtain $z'y'x' = xyz$, Thus $z'y'x'$ and xyz are the same glide reflections, so their axes d, d' as well as vectors v, v' coincide. It follows that lines $d = KL$ and $d' = K'L'$ coincide and $\sigma(T) = |v| = |v'| = \sigma(T')$. This completes the proof.

The idea to apply the same transformation appeared earlier in Vladimir Dubrovsky's solution to the following nice problem proposed by Michael de Villiers [6]: *Prove that the intersections of the adjacent perpendicular bisectors of the sides of a hexagon with opposite sides parallel form a parallelo-hexagon, i.e. hexagon with opposite sides parallel and equal.*

References

- [1] H. S. M. Coxeter, *Introduction to Geometry*, 2nd Edition, John Wiley & Sons, Inc. (1969).
- [2] H. S. M. Coxeter, *The product of three reflections*, Quarterly of Applied Mathematics, Vol. 5, No. 2 (July, 1947), pp. 217–222.
- [3] Clayton W. Dodge, *Euclidean Geometry and Transformations*, Dover Publications (2012).
- [4] Victor Thébault, *Solution to Problem 4470*, The American Mathematical Monthly, Vol. 60, No. 4 (Apr., 1953), pp. 268–269.
- [5] Vladimir Dubrovsky, *Yet another proof of the theorem on isogonal conjugation* (in Russian), Kvant 3/2016, p. 39.
- [6] Vladimir Dubrovsky, *Solution to Michael de Villiers' Problem 7741*, Romantics of Geometry Facebook Group, 22 Nov 2021.
www.facebook.com/groups/parmenides52/posts/4566337916813212/

Edmund Puczyowski (1948–2021) was a distinguished mathematician specializing in the theory non-commutative rings. He was called *Lord of the Rings* by his family and friends. In years 1998–2007 he was a chairman of the Main Committee of the Mathematical Olympiad in Poland. During this period he introduced the highest standards in the organization of the Olympiad and has initiated a very successful Junior Mathematical Olympiad in Poland. He infected everyone with passion for mathematics and mathematics education. He was a very friendly person, thinking more about the others than about himself, always ready to help.

Acknowledgement

The author offers cordial thanks to Vladimir Dubrovsky for suggesting a simpler proof of the second part of Theorem 1.3.