

## Junior problems

J625. The rectangular box  $ABCD A' B' C' D'$  has volume 2023 and total area 2550. Given that

$$\frac{1}{AB} + \frac{1}{AD} - \frac{1}{AA'} = \frac{4}{7},$$

find the dimensions of the box.

*Proposed by Adrian Andreescu, Dallas, USA*

*Solution by Jodie Burdick, SUNY Brockport, NY, USA*

Let  $AB = x, AD = y, AA' = z$ .

Thus the volume is  $xyz$  and the total area is  $2(xy + xz + yz)$ .

This means that  $xyz = 2023$  and  $2(xy + xz + yz) = 2550 \iff xy + xz + yz = 1275$  and so

$$\frac{xy + xz + yz}{xyz} = \frac{1275}{2023} \iff \frac{1}{x} + \frac{1}{y} + \frac{1}{z} = \frac{1275}{2023} \iff \frac{1}{x} + \frac{1}{y} = \frac{1275}{2023} - \frac{1}{z}$$

We also know that

$$\frac{1}{x} + \frac{1}{y} - \frac{1}{z} = \frac{4}{7} \iff \frac{1}{x} + \frac{1}{y} = \frac{4}{7} + \frac{1}{z}$$

Hence

$$\frac{4}{7} + \frac{1}{z} = \frac{1275}{2023} - \frac{1}{z} \iff \frac{2}{z} = \frac{1275}{2023} - \frac{4}{7} \iff \frac{2}{z} = \frac{1}{17} \iff z = 34$$

$xyz = 2023$  implies that

$$xy = \frac{2023}{34} = \frac{119}{2}$$

Moreover

$$\frac{1}{x} + \frac{1}{y} = \frac{4}{7} + \frac{1}{z} \iff \frac{x+y}{xy} = \frac{143}{238} \iff x+y = \frac{143}{238} xy = \frac{143}{238} \frac{119}{2} = \frac{143}{4}$$

Hence  $x$  and  $y$  are the solutions of the quadratic equation

$$t^2 - \frac{143}{4}t + \frac{119}{2} = 0 \iff 4t^2 - 143t + 238 = 0$$

with solutions  $t_1 = 34, t_2 = 7/4$ .

Therefore the solutions of the problem are

$$x = 34, \quad y = 7/4, \quad z = 34$$

and

$$x = 7/4, \quad y = 34, \quad z = 34$$

*Also solved by Soham Dutta, West Bengal, India; Jimi Nguyen, Brockport College, Brockport NY, USA; Matthew Too, Brockport, NY, USA; Vishwesh Ravi Shrimali, Jaipur, India; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Kay Yeon Lee, Choate Rosemary Hall, Wallingford, CT, USA; Austin Lee, The Lawrenceville School, NJ, USA; Soham Bhadra, India; Daniel Pascuas, Barcelona, Spain; Theo Koupelis, Clark College, WA, USA.*

J626. Let  $a, b, c, d$  be positive real numbers. Prove that

$$\frac{bcd}{a} + \frac{acd}{b} + \frac{abd}{c} + \frac{abc}{d} \geq 2\sqrt{a^2b^2 + b^2c^2 + c^2d^2 + d^2a^2}$$

*Proposed by An Zhenping, Xianyang Normal University, China*

*Solution by Daniel Pascuas, Barcelona, Spain*

Since  $\frac{bcd}{a} + \frac{acd}{b} + \frac{abd}{c} + \frac{abc}{d} = abcd\left(\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + \frac{1}{d^2}\right)$ , the inequality can be rewritten as

$$\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2} + \frac{1}{d^2} \geq 2\sqrt{\frac{1}{a^2b^2} + \frac{1}{b^2c^2} + \frac{1}{c^2d^2} + \frac{1}{d^2a^2}},$$

or equivalently

$$x + y + z + t \geq 2\sqrt{xy + yz + zt + tx} \quad (x, y, z, t > 0),$$

which directly follows from the AM-GM inequality since  $xy + yz + zt + tx = (x + z)(y + t)$ .

*Also solved by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy; Soham Dutta, West Bengal, India; Konstantinos Nakis, Athens, Greece; Henry Ricardo, Westchester Area Math Circle; Arkady Alt, San Jose, CA, USA; Jiang Lianjun, Quanzhou Middle School, Guilin, China; CFAL Problem Solving Group, Centre for Advanced Learning, Mangalore, India; Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia; Marian Ursărescu, Romania; Kay Yeon Lee, Choate Rosemary Hall, Wallingford, CT, USA; Austin Lee, The Lawrenceville School, NJ, USA; Soham Bhadra, India; Alexis Norcross, SUNY Brockport, NY, USA; Telemachus Baltsavias, Kerameies Junior High School, Kefalonia, Greece; Ángel Plaza, Universidad de Las Palmas de Gran Canaria, Spain; Sundaresh H R, Shivamogga, India; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Brian Bradie, Christopher Newport University, Newport News, VA, USA; Theo Koupelis, Clark College, WA, USA.*

J627. Let  $a, b, c$  be positive real numbers such that  $a + b + c = 6$ . Prove that

$$\frac{ab}{\sqrt{a^2 + 3a + 6}} + \frac{bc}{\sqrt{b^2 + 3b + 6}} + \frac{ca}{\sqrt{c^2 + 3c + 6}} \leq 3.$$

*Proposed by Mihaela Berindeanu, Bucharest, România*

*Solution by Brian Bradie, Christopher Newport University, Newport News, VA, USA*

Let

$$f(x) = \frac{x}{\sqrt{x^2 + 3x + 6}}.$$

Then

$$f'(x) = \frac{\frac{3}{2}x + 6}{(x^2 + 3x + 6)^{3/2}} \quad \text{and} \quad f''(x) = -\frac{3x^2 + \frac{81}{4}x + 18}{(x^2 + 3x + 6)^{5/2}},$$

so  $f$  is increasing and concave down for  $x > 0$ . By the weighted Jensen's inequality,

$$\begin{aligned} \frac{ab}{\sqrt{a^2 + 3a + 6}} + \frac{bc}{\sqrt{b^2 + 3b + 6}} + \frac{ca}{\sqrt{c^2 + 3c + 6}} &= bf(a) + cf(b) + af(c) \\ &\leq 6f\left(\frac{ab + bc + ca}{6}\right). \end{aligned}$$

Now, adding  $3(ab + bc + ca)$  to both sides of and rearranging terms in the inequality

$$0 \leq \frac{1}{2}(a - b)^2 + \frac{1}{2}(b - c)^2 + \frac{1}{2}(c - a)^2 = a^2 + b^2 + c^2 - ab - bc - ca$$

yields

$$ab + bc + ca \leq \frac{(a + b + c)^2}{3}.$$

Thus,

$$\frac{ab}{\sqrt{a^2 + 3a + 6}} + \frac{bc}{\sqrt{b^2 + 3b + 6}} + \frac{ca}{\sqrt{c^2 + 3c + 6}} \leq 6f\left(\frac{(a + b + c)^2}{18}\right) = 6f(2) = 3.$$

*Also solved by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy; Arkady Alt, San Jose, CA, USA; Jiang Lianjun, Quanzhou Middle School, Guilin, China; Kay Yeon Lee, Choate Rosemary Hall, Wallingford, CT, USA; Soham Dutta, India; Alexander Lee; Telemachus Baltsavias, Kerameies Junior High School, Kefalonia, Greece; Marin Chirciu, Colegiul National Zinca Golescu, Pitesti, Romania; Suvankar Saha, Indian Statistical Institute, India; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Batakogias Panagiotis, High School of Velestino, Greece; Henry Ricardo, Westchester Area Math Circle; Ángel Plaza, Universidad de Las Palmas de Gran Canaria, Spain; Theo Koupelis, Clark College, WA, USA.*

J628. Find all positive integers  $n$  for which

$$(n + 3)! - n! + 7n^3 + 2023$$

is a cube of a prime number.

*Proposed by Adrian Andreescu, Dallas, USA*

*Solution by CFAL Problem Solving Group, Centre for Advanced Learning, Mangalore, India*

Let

$$S(n) = (n + 3)! - n! + 7n^3 + 2023.$$

Since 7 is a factor of 2023, if  $n \geq 7$ , then  $S(n)$  is a multiple of 7. The only prime cube that is a multiple of 7 is  $7^3$ . But

$$[(n + 3)! - n!] + 7n^3 + 2023 > 343$$

for  $n \geq 7$  since the first two terms (i.e.  $[(n + 3)! - n!], 7n^3$ ) are non-negative.

Thus we must have  $n < 7$ . If we try the numbers  $n = 1, 2, 3, 4, 5, 6$ , we see that only  $n = 2$  works for which the prime is 13.

*Also solved by Soham Dutta, West Bengal, India; Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia; Kay Yeon Lee, Choate Rosemary Hall, Wallingford, CT, USA; Austin Lee, The Lawrenceville School, NJ, USA; Soham Bhadra, India; Marin Chirciu, Colegiul National Zinca Golescu, Pitesti, Romania; Sundaresh H R, Shivamogga, India; Theo Koupelis, Clark College, WA, USA.*

J629. Let  $a, b, c$  be positive real numbers such that  $ab + bc + ca = 1$ . Prove that

$$bc\sqrt{2a+b+c} + ca\sqrt{2b+c+a} + ab\sqrt{2c+a+b} \geq \frac{2}{\sqrt{a+b+c}}.$$

*Proposed by Nguyen Viet Hung, Hanoi University of Science, Vietnam*

*Solution by the author*

First, we have

$$\left( \sum_{\text{cyc}} bc\sqrt{2a+b+c} \right)^2 \left( \sum_{\text{cyc}} \frac{bc}{\sqrt{2a+b+c}} \right) \geq (bc + ca + ab)^3 = 1.$$

On the other hand, the Cauchy-Schwarz inequality also yields

$$\begin{aligned} \sum_{\text{cyc}} \frac{bc}{\sqrt{2a+b+c}} &\leq (bc + ca + ab) \left( \sum_{\text{cyc}} \frac{bc}{2a+b+c} \right) \\ &= \sum_{\text{cyc}} \frac{bc}{(a+b) + (a+c)} \\ &\leq \sum_{\text{cyc}} \frac{bc}{4} \left( \frac{1}{a+b} + \frac{1}{a+c} \right) \\ &= \frac{a+b+c}{4}. \end{aligned}$$

Combining these two relations we get the desired inequality.

*Also solved by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy; Batakogias Panagiotis, High School of Velestino, Greece; Jiang Lianjun, Quanzhou Middle School, Guilin, China; Marin Chirciu, Colegiul National Zinca Golescu, Pitesti, Romania; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Ángel Plaza, Universidad de Las Palmas de Gran Canaria, Spain; Theo Koupelis, Clark College, WA, USA.*

J630. Let  $a, b, c$  be positive real numbers such that  $abc = 1$ . Prove that

$$\frac{7}{a} + \frac{7}{b} + \frac{7}{c} + \frac{16}{a+b} + \frac{16}{b+c} + \frac{16}{c+a} + \frac{27}{a+b+c} \geq 54.$$

*Proposed by Marius Stănean, Zalău, România*

*Solution by Theo Koupelis, Clark College, WA, USA*

Without loss of generality let  $a \geq b \geq c$ , and

$$f(a, b, c) := \frac{7}{a} + \frac{7}{b} + \frac{7}{c} + \frac{16}{a+b} + \frac{16}{b+c} + \frac{16}{c+a} + \frac{27}{a+b+c}.$$

For  $b = c$  and  $ab^2 = 1$  we get

$$\begin{aligned} f(a, b, b) &= \frac{7}{a} + \frac{22}{b} + \frac{32}{a+b} + \frac{27}{a+2b} = 7b^2 + \frac{22}{b} + \frac{32b^2}{1+b^3} + \frac{27b^2}{1+2b^3} \\ &= \frac{2(7b^9 + 78b^6 + 66b^3 + 11)}{b(1+b^3)(1+2b^3)}, \end{aligned}$$

and thus

$$f(a, b, b) \geq 54 \iff 2(b-1)^4(7b^5 + 28b^4 + 16b^3 + 2b^2 + 17b + 11) \geq 0,$$

which is obvious. Also,

$$f(a, b, c) - f(a, b, b) = (b-c) \cdot \left[ \frac{7}{bc} + \frac{16}{2b(b+c)} + \frac{16}{(a+b)(a+c)} + \frac{27}{(a+b+c)(a+2b)} \right] \geq 0.$$

Thus,  $f(a, b, c) \geq 54$ , with equality when  $a = b = c = 1$ .

*Also solved by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy; Austin Lee, The Lawrenceville School, NJ, USA; Arkady Alt, San Jose, CA, USA; Telemachus Baltasavias, Kerameies Junior High School, Kefalonia, Greece; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

## Senior problems

S625. Prove that there are infinitely many positive integers  $n$  such that  $n^2 + 5$  has a proper divisor greater than  $8n/5$ .

*Proposed by Titu Andreescu, University of Texas at Dallas, USA*

*First solution by the author*

Use the identity

$$(L_m)^2 - L_{m-1}L_{m+1} = 5(-1)^{m-1},$$

where  $(L_m)_{m \geq 0}$  is the Lucas sequence:  $L_0 = 2, L_1 = 1, L_{m+1} = L_m + L_{m-1}, m = 1, 2, 3, \dots$  to get

$$(L_{2k})^2 + 5 = L_{2k-1}L_{2k+1}.$$

Take  $n = L_{2k}$  and use the fact that  $L_{m+1} > \frac{8}{5}L_m$ , for  $m \geq 5$ , which can be proven by an easy strong induction.

*Second solution by Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia*

Let's call  $n \in \mathbb{N}$  good if  $n^2 + 5$  has a proper divisor greater than  $\frac{8n}{5}$ . We will show that all odd positive integers are good.

If  $n \in \{1, 3\}$  then  $\frac{n^2+5}{2}$ , which is the greatest proper divisor of  $n^2 + 5$ , is clearly greater than  $\frac{8n}{5}$ .

If odd  $n \geq 5$ , let  $n = 2k + 1, k \geq 2$ . We still consider  $\frac{n^2+5}{2} = 2k^2 + 2k + 3$  as the greatest proper divisor of  $n^2 + 5$ . In fact,  $\forall k \geq 2$ , we have

$$\frac{8n}{5} = \frac{16k+8}{5} \leq 4k < 4k+3+2k(k-1) = \frac{n^2+5}{2}.$$

The proof is complete.

*Also solved by Soham Dutta, West Bengal, India; Kay Yeon Lee, Choate Rosemary Hall, Wallingford, CT, USA; Austin Lee, The Lawrenceville School, NJ, USA; Brian Bradie, Christopher Newport University, Newport News, VA, USA; Theo Koupelis, Clark College, WA, USA; Telemachus Baltsavias, Kerameies Junior High School, Kefalonia, Greece; Chayan Kumar Mandal, Malda, West Bengal, India; Sundaresh H R, Shivamogga, India.*

S626. Let  $a, b, c, d, e$  be nonnegative real numbers such that  $ab + bc + cd + de + ea = 1$ . Prove that

$$3 < \frac{1}{a+1} + \frac{1}{b+1} + \frac{1}{c+1} + \frac{1}{d+1} + \frac{1}{e+1} \leq 4.$$

*Proposed by Vasile Cîrtoaje, Oil-Gas University, Ploiești, România*

*Solution by the author*

Let  $(x_1, x_2, x_3, x_4, x_5)$  be a permutation of  $(a, b, c, d, e)$  such that  $x_1 \geq x_2 \geq x_3 \geq x_4 \geq x_5$ . Due to symmetry, the desired inequality is equivalent to

$$\frac{1}{x_1+1} + \frac{1}{x_2+1} + \frac{1}{x_3+1} + \frac{1}{x_4+1} + \frac{1}{x_5+1} \leq 4,$$

that can be written in the form

$$\frac{x_1}{x_1+1} + \frac{x_2}{x_2+1} + \frac{x_3}{x_3+1} + \frac{x_4}{x_4+1} + \frac{x_5}{x_5+1} \geq 1.$$

For  $x_1x_2 \geq 1$ , the inequality is true because

$$\frac{x_1}{x_1+1} + \frac{x_2}{x_2+1} - 1 = \frac{x_1x_2 - 1}{(x_1+1)(x_2+1)} \geq 0.$$

Consider further  $x_1x_2 \leq 1$ . For  $x_3 + x_4 + x_5 > 0$ , by the Cauchy-Schwarz inequality we have

$$\begin{aligned} \frac{x_3}{x_3+1} + \frac{x_4}{x_4+1} + \frac{x_5}{x_5+1} &\geq \frac{(x_3 + x_4 + x_5)^2}{x_3(x_3+1) + x_4(x_4+1) + x_5(x_5+1)} \\ &\geq \frac{(x_3 + x_4 + x_5)^2}{(x_3 + x_4 + x_5)^2 + x_3 + x_4 + x_5} = \frac{x_3 + x_4 + x_5}{x_3 + x_4 + x_5 + 1} = 1 - \frac{1}{x_3 + x_4 + x_5 + 1}, \end{aligned}$$

hence

$$\frac{x_3}{x_3+1} + \frac{x_4}{x_4+1} + \frac{x_5}{x_5+1} \geq 1 - \frac{1}{x_3 + x_4 + x_5 + 1}.$$

We can see that this inequality is also true for  $x_3 = x_4 = x_5 = 0$ . So, it suffices to prove that

$$\frac{x_1}{x_1+1} + \frac{x_2}{x_2+1} - \frac{1}{x_3 + x_4 + x_5 + 1} \geq 0.$$

By Lemma below, we have

$$x_3 + x_4 + x_5 \geq \frac{1 - x_1x_2}{x_1 + x_2},$$

hence

$$\begin{aligned} \frac{x_1}{x_1+1} + \frac{x_2}{x_2+1} - \frac{1}{x_3 + x_4 + x_5 + 1} &\geq \frac{x_1}{x_1+1} + \frac{x_2}{x_2+1} - \frac{1}{\frac{1-x_1x_2}{x_1+x_2} + 1} \\ &= \frac{x_1}{x_1+1} + \frac{x_2}{x_2+1} - \frac{x_1+x_2}{1-x_1x_2+x_1+x_2} = \frac{2x_1x_2(1-x_1x_2)}{(x_1+1)(x_2+1)(1-x_1x_2+x_1+x_2)} \geq 0. \end{aligned}$$

The proof is completed. The equality is an equality for  $ab = 1$  and  $c = d = e = 0$  (or any cyclic permutation).



*Lemma:* Let  $a, b, c, d, e$  be nonnegative real numbers, and let  $(x_1, x_2, x_3, x_4, x_5)$  be a permutation of  $(a, b, c, d, e)$  such that  $x_1 \geq x_2 \geq x_3 \geq x_4 \geq x_5$ . Then,

$$x_1x_2 + (x_1 + x_2)(x_3 + x_4 + x_5) \geq ab + bc + cd + de + ea.$$

*Proof:* Assume that  $a = \max\{a, b, c, d, e\}$ , hence  $a = x_1$  and  $x_2 + x_3 + x_4 + x_5 = b + c + d + e$ . Since

$$\begin{aligned} x_1x_2 + (x_1 + x_2)(x_3 + x_4 + x_5) &= x_1(x_2 + x_3 + x_4 + x_5) + x_2(x_3 + x_4 + x_5) \\ &= a(b + c + d + e) + x_2(x_3 + x_4 + x_5) \geq a(b + c + d + e) + x_2x_3, \end{aligned}$$

it suffices to show that

$$a(b + c + d + e) + x_2x_3 \geq ab + bc + cd + de + ea,$$

that is

$$a(c + d) + x_2x_3 \geq bc + cd + de,$$

which is equivalent to the obvious inequality

$$c(a - b) + d(a - c) + (x_2x_3 - de) \geq 0.$$

*Also solved by Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

S627. Let  $a, b, c$  be positive real numbers. Prove that

$$2a\sqrt{9b^2 + 16c^2} + 2b\sqrt{9c^2 + 16a^2} + 2c\sqrt{9a^2 + 16b^2} + 15abc \left( \frac{1}{2b + 3c} + \frac{1}{2c + 3a} + \frac{1}{2a + 3b} \right) \geq 13(ab + bc + ca)$$

*Proposed by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy*

*Solution by the author*

$$\sqrt{9b^2 + 16c^2} \geq 3b + 4c - \frac{10bc}{2b + 3c} \iff \frac{24bc(b - c)^2}{(2b + 3c)^2} \geq 0$$

hence we come to prove

$$2 \sum_{\text{cyc}} (3ab + 4ac) - 20abc \sum_{\text{cyc}} \frac{1}{2b + 3c} + 15abc \sum_{\text{cyc}} \frac{1}{2b + 3c} \geq 13(ab + bc + ca)$$

that is

$$ab + bc + ca \geq 5abc \left( \frac{1}{2b + 3c} + \frac{1}{2c + 3a} + \frac{1}{2a + 3b} \right)$$

The AGM yields

$$ab + bc + ca \geq 5abc \left( \frac{1}{5b^{2/5}c^{3/5}} + \frac{1}{5c^{2/5}a^{3/5}} + \frac{1}{5a^{2/5}b^{3/5}} \right) = b^{3/5}c^{2/5}a + c^{3/5}a^{2/5}b + a^{3/5}b^{2/5}c$$

and the AGM again finishes the proof

$$(ab + ab + ab + ac + ac)/5 \geq ab^{3/5}c^{2/5} \text{ and cyclic}$$

*Also solved by Theo Koupelis, Clark College, WA, USA; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

S628. Prove that there are infinitely many positive integers  $n$  such that precisely two of the numbers  $n - 2$ ,  $n + 2$ ,  $5(n - 2)$ ,  $5(n + 2)$  are perfect squares.

*Proposed by Titu Andreescu, University of Texas at Dallas, USA*

*Solution by Brian Bradie, Christopher Newport University, Newport News, VA, USA*

We start with two observations. For  $n > 2$ ,

- $n - 2$  and  $n + 2$  cannot simultaneously be perfect squares; and
- if  $n - 2$  is a perfect square, then  $5(n - 2)$  cannot be a perfect square as well.

Thus, it suffices to show that there are infinitely many positive integers  $n$  such that  $n - 2$  and  $5(n + 2)$  are perfect squares. Suppose  $n - 2 = m^2$  and  $5(n + 2) = \ell^2$  for positive integers  $m$  and  $\ell$ . This leads to the Pell equation  $\ell^2 - 5m^2 = 20$ , which has solutions

$$(\ell_k, m_k) = (5F_{2k-1}, L_{2k-1})$$

for  $k = 1, 2, 3, \dots$ , where  $F_k$  and  $L_k$  are the  $k$ th Fibonacci number and the  $k$ th Lucas number, respectively. Thus, for  $n = L_{2k-1}^2 + 2$  with  $k = 1, 2, 3, \dots$ ,  $n - 2$  and  $5(n + 2)$  are perfect squares.

*Also solved by Soham Dutta, West Bengal, India; Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia; Theo Koupelis, Clark College, WA, USA; Telemachus Baltsavias, Kerameies Junior High School, Kefalonia, Greece; Batakogias Panagiotis, High School of Velestino, Greece; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

S629. Let  $ABCD$  be a rectangle with area  $[ABCD]$  and  $O$  a point in its plane. Prove that

$$|OA \cdot OC - OB \cdot OD| \leq [ABCD] \leq OA \cdot OC + OB \cdot OD.$$

*Proposed by Jozsef Tkadlec, Czech Republic*

*Solution by Theo Koupelis, Clark College, WA, USA*

Let the center  $K$  of the rectangle be at the origin of a Cartesian coordinate system so that  $A = (-a, b)$ ,  $B = (-a, -b)$ ,  $C = (a, -b)$ , and  $D = (a, b)$ , where  $a, b > 0$ . Then  $[ABCD] = 4ab$ . If  $O = (x, y)$ , then using Cauchy-Schwarz we get

$$OA \cdot OC = \sqrt{(x+a)^2 + (y-b)^2} \sqrt{(x-a)^2 + (y+b)^2} \geq |(x+a)(y+b)| + |(x-a)(y-b)|,$$

and

$$OB \cdot OD = \sqrt{(x+a)^2 + (y+b)^2} \sqrt{(x-a)^2 + (y-b)^2} \geq |(x+a)(y-b)| + |(x-a)(y+b)|.$$

Thus,

$$OA \cdot OC + OB \cdot OD \geq (|x+a| + |x-a|)(|y+b| + |y-b|)$$

Setting  $f(x) = |x+a| + |x-a|$ , we easily find that  $f(x) = 2|x|$  when  $|x| \geq a$ , and  $f(x) = 2a$  when  $|x| < a$ . Similarly, setting  $g(y) = |y+b| + |y-b|$ , we find  $g(y) = 2|y|$  when  $|y| \geq b$ , and  $g(y) = 2b$  when  $|y| < b$ . Thus, in all cases  $OA \cdot OC + OB \cdot OD \geq 4ab = [ABCD]$ .

The inequality on the left-hand-side is equivalent to

$$\begin{aligned} OA \cdot OC + OB \cdot OD &\geq \frac{|OA^2 \cdot OC^2 - OB^2 \cdot OD^2|}{4ab} \iff \\ OA \cdot OC + OB \cdot OD &\geq \frac{|16abxy|}{4ab} = 4|xy|, \end{aligned}$$

which is obvious from the above analysis of the behavior of the functions  $f(x)$  and  $g(y)$ . Both equalities occur when  $O$  is one of the vertices of the rectangle. The equality on the right-hand-side also occurs when  $O = K$  in the case when  $a = b$ .

*Also solved by Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

S630. Let  $n \geq 2$  and  $x_1, \dots, x_n$  be real numbers, not all zero, adding up to zero. Moreover, for each positive real number  $t$  there are at most  $\frac{1}{t}$  pairs  $(i, j)$  such that  $|x_i - x_j| \geq t$ . Prove that

$$x_1^2 + \dots + x_n^2 < \frac{1}{n} \left( \max_{1 \leq i \leq n} x_i - \min_{1 \leq i \leq n} x_i \right).$$

*Proposed by Navid Safaei, Sharif University of Technology, Tehran, Iran*

*Solution by the author*

Notice that  $n \sum_{i=1}^n x_i^2 = (\sum_{i=1}^n x_i)^2 + \sum_{1 \leq i < j \leq n} (x_i - x_j)^2$ . It then suffices to prove

$$\sum_{1 \leq i < j \leq n} (x_i - x_j)^2 \leq \max_{1 \leq i \leq n} x_i - \min_{1 \leq i \leq n} x_i.$$

Let us denote  $\{|x_i - x_j| : 1 \leq i < j \leq n\} = \{y_1, \dots, y_m\}$ , then assume  $y_1 < y_2 < \dots < y_m$ . Then assume that the total number of pairs  $(i, j)$  such that  $|x_i - x_j| = y_k$  is  $a_k$ . Letting  $t = y_k$  then there are at most  $\frac{1}{y_k}$  pairs  $(i, j)$  such that  $|x_i - x_j| \geq t$ . Hence,

$$a_k + a_{k+1} + \dots + a_m \leq \frac{1}{y_k}.$$

Letting  $S_k = a_k + a_{k+1} + \dots + a_m$  it follows that  $S_k \leq \frac{1}{y_k}$ .

Hence,

$$\sum_{1 \leq i < j \leq n} (x_i - x_j)^2 = \frac{1}{2} \sum_{k=1}^m a_k y_k^2 = \frac{1}{2} \sum_{k=1}^m S_k (y_k^2 - y_{k-1}^2) \leq \frac{1}{2} \sum_{k=1}^m \frac{y_k^2 - y_{k-1}^2}{y_k}$$

Putting  $y_0 = 0$  and  $y_{k-1} < y_k$  it follows that  $\frac{y_k^2 - y_{k-1}^2}{y_k} < 2(y_k - y_{k-1})$ . Hence,

$$\sum_{1 \leq i < j \leq n} (x_i - x_j)^2 < 2 \sum_{k=1}^m (y_k - y_{k-1}) = y_m = \max_{1 \leq i \leq n} x_i - \min_{1 \leq i \leq n} x_i.$$

*Also solved by Daniel Pascuas, Barcelona, Spain.*

## Undergraduate problems

U625. Evaluate

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{\pi \sin \frac{\pi k}{2n}}{2n(\sin \frac{\pi k}{2n} + \cos \frac{\pi k}{2n})}.$$

*Proposed by Ángel Plaza, University of Las Palmas de Gran Canaria, Spain*

*Solution by Henry Ricardo, Westchester Area Math Circle*

Rewriting the given summation, we recognize it as a right Riemann sum so that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\pi}{2} \cdot \sum_{k=1}^n \frac{\sin \frac{\pi}{2} \cdot \frac{k}{n}}{\sin \frac{\pi}{2} \cdot \frac{k}{n} + \cos \frac{\pi}{2} \cdot \frac{k}{n}} \cdot \frac{1}{n} &= \frac{\pi}{2} \int_0^1 \frac{\sin \frac{\pi x}{2}}{\sin \frac{\pi x}{2} + \cos \frac{\pi x}{2}} dx \\ &\stackrel{u=\pi x/2}{=} \int_0^{\pi/2} \frac{\sin u}{\sin u + \cos u} du \\ &= \frac{1}{2} \left( \int_0^{\pi/2} \frac{\sin u + \cos u}{\sin u + \cos u} du - \int_0^{\pi/2} \frac{\cos u - \sin u}{\sin u + \cos u} du \right) \\ &= \frac{\pi}{4}. \end{aligned}$$

*Also solved by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy; Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia; Daniel Pascuas, Barcelona, Spain; Brian Bradie, Christopher Newport University, Newport News, VA, USA; Theo Koupelis, Clark College, WA, USA; Seán M. Stewart, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia; Etisha Sharma, Agra College, Agra, India; Matthew Too, Brockport, NY, USA; Soham Dutta, India; Sundares H R, Shivamogga, India; Suvankar Saha, Indian Statistical Institute, India; Swastika Dey, India; Toyesh Prakash Sharma, Agra College, India; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Prajnanaswaroop S, Bengaluru, Karnataka, India.*

U626. Let

$$P(x) = 99x^6 + 3x^5 + x^4 + 2x^3 + 4x^2 - 1.$$

Prove that there is a prime  $q$  such that  $P(q) = (q - 1)!$ .

*Proposed by Titu Andreescu, University of Texas at Dallas, USA*

*Solution by Theo Koupelis, Clark College, WA, USA*

For  $q \geq 19$  we have  $(q - 1)! = 5! \cdot 6 \cdot 7 \cdots (q - 2)(q - 1) > 100q^6$ , because  $5! > 100$  and  $k(q - k + 1) > q \Leftrightarrow q > k$ , where  $k \in \{6, 7, 8, 9, 10, 11\}$ . But for any prime  $p \geq 11$  we have  $99p^6 + 3p^5 + p^4 + 2p^3 + 4p^2 - 1 < 100p^6$  because  $p^6 \geq 11p^5 > (3+1+2+4)p^5 > 3p^5 + p^4 + 2p^3 + 4p^2 - 1$ . Therefore, there is no solution for  $q \geq 19$ . On the other hand, it is clear that  $P(x) > 99x^6$  for any positive integer, and thus  $P(q) > 99 \cdot 2^6 = 6336$ , while  $(8 - 1)! = 7! = 5040$ . Therefore, there is no solution for  $q \leq 7$ . Thus, we only need examine the cases  $q = 11, 13, 17$ . By substituting, we find that  $P(q) = (q - 1)!$  only for  $q = 13$ .

*Also solved by Soham Dutta, West Bengal, India; Sophia Tiffany, SUNY Brockport, NY, USA; Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia; Soham Bhadra, India; Juan José Granier, Universidad de Chile, Chile; Prajnanaswaroop S, Bengaluru, Karnataka, India; Telemachus Baltasavias, Kerameies Junior High School, Kefalonia, Greece; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

U627. The author claims that he had a nice solution but the margin was too small :)



U628. Let  $Q(x, y)$  be the set of rational functions with rational coefficients and  $S$  be the set of symmetric rational functions with rational coefficients. Find all  $a, b$  such that

$$Q(x^a + y^a, x^b + y^b) = S$$

*Proposed by Navid Safaei, Sharif University of Technology, Tehran, Iran*

*Solution by the author*

Let  $N_a = x^a + y^a$ . Notice that if  $P_1, P_2$  are members of  $Q(N_a, N_b)$  then  $\frac{P_1}{P_2}, P_1 + P_2, P_i^k$  are members of  $Q(N_a, N_b)$ . Moreover, all symmetric expressions in two variables can be written as a rational function of  $x + y, xy$ . So, it suffices to check whether or not  $x + y, xy$  can be constructed or not.

Indeed,  $S_1 = x + y, S_2 = xy$ . Notice that  $S_2 = \frac{N_1^2 - N_2}{2}$ . So,  $Q(N_2, N_1) = S$ . Further, since  $N_1^3 - N_3 = 3N_1S_2$  implies  $S_2 = \frac{N_1^3 - N_3}{3N_1}$ . Thus,  $Q(N_1, N_3) = S$ .

Now, assume  $b \geq a \geq 2$ . Assume  $Q(N_a, N_b) = S$  thus,  $x + y = S$ . Hence, there must be coprime polynomials  $P_1, P_2$  such that

$$\frac{P_1(x^a + y^a, x^b + y^b)}{P_2(x^a + y^a, x^b + y^b)} = x + y$$

If  $\gcd(a, b) = d > 1$  then the map  $x \rightarrow \omega x$  where  $\omega$  is the primitive  $d$ -th root of unity preserves the left while changes the right. Hence, we can assume that  $\gcd(a, b) = 1$ . Let  $\xi$  be the primitive  $a$ -th root of  $-1$ ;  $\xi^a = -1$  then the map  $y \rightarrow \xi x$  yields;

$$P_1(0, (\xi^b + 1)x^b) = (\xi + 1)xP_2(0, (\xi^b + 1)x^b)$$

Since  $a \neq 1, \gcd(a, b) = 1$  it follows that  $\xi + 1, \xi^b + 1 \neq 0$ . Let  $\alpha$  be the primitive  $b$ -th root of unity, that is  $\alpha^b = 1$  the map  $x \rightarrow \alpha x$  preserves everything except  $x$ . It follows that for each  $x$ ;

$$P_1(0, (\xi^b + 1)x^b) = P_2(0, (\xi^b + 1)x^b) = 0.$$

But then,  $P_1, P_2$  will have common factors. This is impossible. Thus,  $x + y$  can not be represented as such.

Finally, if  $a = 1, b \geq 4$  we shall prove that  $xy$  can not be constructed. On the contrary, assume there are coprime polynomials  $P_1, P_2$  such that

$$\frac{P_1(x + y, x^b + y^b)}{P_2(x + y, x^b + y^b)} = xy.$$

That is,  $P_1(x + y, x^b + y^b) = xyP_2(x + y, x^b + y^b)$ . Let  $\beta$  be some of  $b$ -th root of  $-1$ , that is,  $\beta^b = -1$  such that  $\beta \neq -1$ .

Then, the map  $y \rightarrow \beta x$  leads to the following expression

$$P_1((\beta + 1)x, 0) = \beta x^2 P_2((\beta + 1)x, 0).$$

That is,  $P_1(x, 0) = \frac{\beta}{(\beta + 1)^2} x^2 P_2(x, 0)$ . This means that for all  $b - 1 \geq 3$  roots of  $z^b = -1$  such that  $z \neq -1$  the fraction  $\frac{\beta}{(\beta + 1)^2}$  must be constant. But then, the equation  $\frac{x}{(x + 1)^2} = C$  must have  $b - 1$  roots, however, it can at most have two distinct roots. This is a contradiction.

U629. Let  $D = \{z \in \mathbb{C} : |z| < 1\}$  and let  $f : D \rightarrow \mathbb{C}$  be a holomorphic function such that  $|f(z)| \leq 1$  and  $f(0) = 0$ . Prove that for  $z \neq 0$  the following inequality holds

$$\frac{|f(z)|}{|z|(1 + |f'(0)|)} + \frac{|z||f'(0)|}{|z| + |f(z)|} \leq 1.$$

*Proposed by Alessandro Ventullo, Milan, Italy*

*Solution by Theo Koupelis, Clark College, WA, USA*

Clearing denominators, the desired inequality is equivalent to

$$|f(z)|^2 - |f(z)| \cdot |z||f'(0)| - |z|^2(1 - |f'(0)|^2) \leq 0.$$

Thus, solving the quadratic in  $|f(z)|$ , it is sufficient to show that

$$\frac{1}{2} \left( |z||f'(0)| - |z|\sqrt{4 - 3|f'(0)|^2} \right) \leq |f(z)| \leq \frac{1}{2} \left( |z||f'(0)| + |z|\sqrt{4 - 3|f'(0)|^2} \right).$$

But from the given conditions and the Schwarz lemma we get that  $|f(z)| \leq |z|$  for all  $z \in \mathbb{D}$ , and  $|f'(0)| \leq 1$ . Clearly then  $|z||f'(0)| \leq |z|\sqrt{4 - 3|f'(0)|^2}$ , and thus, it is sufficient to show that

$$|z| \leq \frac{1}{2} \left( |z||f'(0)| + |z|\sqrt{4 - 3|f'(0)|^2} \right),$$

or, for  $z \neq 0$ ,

$$2 - |f'(0)| \leq \sqrt{4 - 3|f'(0)|^2}.$$

After squaring and simplifying we get  $|f'(0)|^2 \leq |f'(0)|$ , which is obvious.

*Also solved by Prajnanaswaroop S, Bengaluru, Karnataka, India.*

U630. Evaluate

$$\int_{-\infty}^{+\infty} \frac{(\tanh y)(\sinh y)^2}{2+(\sinh(2y))^2} y^3 dy$$

*Proposed by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy*

*Solution by the author*

The integral becomes

$$\int_{-1}^1 \frac{x \frac{x^2}{1-x^2}}{2 + \frac{4x^2}{(1-x^2)^2}} \frac{1}{8} \left( \ln \frac{1+x}{1-x} \right)^3 \frac{1}{1-x^2} dx = \frac{1}{16} \int_{-1}^1 \frac{x^3}{1+x^4} \left( \ln \frac{1+x}{1-x} \right)^3 dx$$

We employ the residue–theory. Define the complex function

$$\frac{z^3}{1+z^4} \left( \operatorname{Ln} \frac{1+z}{1-z} \right)^4 \doteq g(z)f^4(z)$$

defined over all  $\mathbf{C}$  cut on the interval  $[-1, 1]$  where  $\operatorname{Ln}(\rho e^{it}) = \ln \rho + it$  (we reserve the notation  $\ln y$  for real arguments  $y$ ). The path of integration runs counterclockwise and encircles  $[-1, 1]$ , close to it. When the maximum distance between the path and the interval  $[-1, 1]$  goes to zero, the integral becomes

$$\int_1^{-1} g(x)f^4(x)dx + \int_{-1}^1 g(x)(f(x) + 2\pi i)^4 dx = -2\pi i \sum (\operatorname{Res}(g(z)f^4(z)))$$

The residues are calculated at the points  $z_k = e^{i\frac{\pi}{4} + k\frac{\pi}{2}}$ ,  $k = 0, 1, 2, 3$  and at  $z = \infty$ . The points  $z_k$  are clearly the zeroes of  $z^4 = -1$ . We get

$$\begin{aligned} & \int_1^{-1} g f^4 dx + \int_{-1}^1 g f^4 dx + \int_{-1}^1 g (8\pi i f^3 + 4(2\pi i)^3 f + 6f^2(2\pi i)^2) dx = \\ & = -2\pi i \sum \operatorname{Res}(g(z)f^4(z)) \end{aligned}$$

that is

$$4I = - \sum \operatorname{Res}(g(z)f^4(z)) + 16\pi^2 \int_{-1}^1 g f dx$$

where we have used  $\int_{-1}^1 g dx = \int_{-1}^1 g f^2 dx = 0$  by oddness.

To compute

$$\int_{-1}^1 g(x)f(x) dx$$

we proceed in the same way writing

$$\int_1^{-1} g f^2 dx + \int_{-1}^1 g (f + 2\pi i)^2 dx = -2\pi i \sum \operatorname{Res} g(z)f^2(z)$$

which yields

$$\int_{-1}^1 g f dx = -\frac{1}{2} \sum \operatorname{Res}(g(z)f^2(z))$$

To compute the residue at  $z = \infty$  of the function  $g(z)f^2(z)$  we compute it on the curve  $\gamma(t) \in \mathbf{C}$ ,  $z = \gamma(t) = it$  and obtain

$$\begin{aligned} g(z)f^2(z) \Big|_{z=\gamma(t)} &= \frac{1}{it} \frac{1}{1 + \frac{1}{t^4}} \left( \operatorname{Ln} \frac{\sqrt{1+t^2} e^{i \arctan t}}{\sqrt{1+t^2} e^{-i \arctan t}} \right)^2 = \\ &= \frac{1}{it} \frac{1}{1 + \frac{1}{t^4}} \left( \operatorname{Lne}^{-2i \arctan \frac{1}{t} + i\pi} \right)^2 = \frac{1}{it} \frac{1}{1 + \frac{1}{t^4}} \left( -i \left( \frac{1}{t} + O\left(\frac{1}{t^3}\right) + \pi \right) \right)^2 = \\ &= -\frac{\pi^2}{it} + O(1/t^2) \end{aligned}$$

whence the residue equal to  $\pi^2$ . The residue in the other four points are

$$\lim_{z \rightarrow z_k} g(z)(z - z_k)f^2(z)$$

because the function has simple poles at any point  $z_k$ ,  $k = 0, 1, 2, 3$ . Denoting  $\varphi_0 = \pi/4$  we get

$$\begin{aligned} \lim_{z \rightarrow z_0} g(z)(z - z_0)f^2(z) &= \lim_{z \rightarrow z_0} \frac{3z^2(z - z_0) + z^3}{4z^3} \left( \operatorname{Ln} \frac{1 + z_0}{1 - z_0} \right)^2 = \\ &= \frac{1}{4} \left( \operatorname{Ln} \frac{1 + e^{i\varphi_0}}{1 - e^{i\varphi_0}} \right)^2 = \frac{1}{4} \left( \operatorname{Ln} \frac{i}{\tan \frac{\varphi_0}{2}} \right)^2 = \frac{1}{4} \left( \frac{1}{2}\pi i + \ln(1 + \sqrt{2}) \right)^2 \end{aligned}$$

$\varphi_1 = 3\pi/4$  we get

$$\begin{aligned} \lim_{z \rightarrow z_1} g(z)(z - z_1)f^2(z) &= \lim_{z \rightarrow z_1} \frac{3z^2(z - z_1) + z^3}{4z^3} \left( \operatorname{Ln} \frac{1 + z_1}{1 - z_1} \right)^2 = \\ &= \frac{1}{4} \left( \operatorname{Ln} \frac{1 + e^{i\varphi_1}}{1 - e^{i\varphi_1}} \right)^2 = \frac{1}{4} \left( \operatorname{Ln} \frac{i}{\tan \frac{\varphi_1}{2}} \right)^2 = \frac{1}{4} \left( \frac{1}{2}\pi i + \ln(\sqrt{2} - 1) \right)^2 \end{aligned}$$

$\varphi_2 = 5\pi/4$  we get

$$\begin{aligned} \lim_{z \rightarrow z_2} g(z)(z - z_2)f^2(z) &= \lim_{z \rightarrow z_2} \frac{3z^2(z - z_2) + z^3}{4z^3} \left( \operatorname{Ln} \frac{1 + z_2}{1 - z_2} \right)^2 = \\ &= \frac{1}{4} \left( \operatorname{Ln} \frac{1 + e^{i\varphi_2}}{1 - e^{i\varphi_2}} \right)^2 = \frac{1}{4} \left( \operatorname{Ln} \frac{i}{\tan \frac{\varphi_2}{2}} \right)^2 = \frac{1}{4} \left( \frac{3}{2}\pi i + \ln(\sqrt{2} - 1) \right)^2 \end{aligned}$$

$\varphi_3 = 7\pi/4$  we get

$$\begin{aligned} \lim_{z \rightarrow z_3} g(z)(z - z_3)f^2(z) &= \lim_{z \rightarrow z_3} \frac{3z^2(z - z_3) + z^3}{4z^3} \left( \operatorname{Ln} \frac{1 + z_3}{1 - z_3} \right)^2 = \\ &= \frac{1}{4} \left( \operatorname{Ln} \frac{1 + e^{i\varphi_3}}{1 - e^{i\varphi_3}} \right)^2 = \frac{1}{4} \left( \operatorname{Ln} \frac{i}{\tan \frac{\varphi_3}{2}} \right)^2 = \frac{1}{4} \left( \frac{3}{2}\pi i + \ln(\sqrt{2} + 1) \right)^2 \end{aligned}$$

Their sum is (use  $\ln(1 + \sqrt{2}) = \ln(\sqrt{2} - 1)^{-1}$ )

$$\begin{aligned} &\frac{1}{4} \left( -2\frac{9}{4}\pi^2 - 2\frac{\pi^2}{4} + 3\pi i \ln(\sqrt{2} - 1) + 3\pi i \ln(1 + \sqrt{2}) + \pi i \ln(\sqrt{2} - 1) + \right. \\ &\left. + \pi i \ln(1 + \sqrt{2}) + 2\ln^2(\sqrt{2} - 1) + 2\ln^2(1 + \sqrt{2}) \right) = \\ &= \frac{1}{4} \left( -2\frac{9}{4}\pi^2 - 2\frac{\pi^2}{4} + 2\ln^2(\sqrt{2} - 1) + 2\ln^2(1 + \sqrt{2}) \right) = -\frac{5}{4}\pi^2 + \ln^2(1 + \sqrt{2}) \end{aligned}$$

Finally

$$\begin{aligned} -\frac{1}{2} \sum \operatorname{Res}(g(z)f^2(z)) &= -\frac{\pi^2}{2} - \frac{1}{2} \sum_{k=0}^3 \lim_{z \rightarrow z_k} g(z)(z - z_k)f^2(z) = \\ &= \frac{\pi^2}{8} - \frac{1}{2}(\ln^2(1 + \sqrt{2})) = \int_{-1}^1 g(x)f^2(x)dx \end{aligned}$$

The last step is to compute

$$\sum \operatorname{Res}(g(z)f^4(z)) \quad (*)$$

The residue at  $z = \infty$  follows by

$$g(z)f^4(z)\Big|_{z=\gamma(t)} = \frac{1}{it} \frac{1}{1 + \frac{1}{t^4}} \left( \text{Lne}^{-i\left(\frac{1}{t} + O\left(\frac{1}{t^3}\right) - \pi\right)} \right)^4 = \frac{1}{it} \left(1 + O\left(\frac{1}{t}\right)\right) (\pi^4 + O(1/t))$$

whence the residue is equal to  $-\pi^4$ .

$$\text{Res}(gf^4)\Big|_{z=z_0} = \frac{1}{4} \left( \frac{1}{2} \pi i + \ln(1 + \sqrt{2}) \right)^4, \quad \text{Res}(gf^4)\Big|_{z=z_1} = \frac{1}{4} \left( \frac{1}{2} \pi i + \ln(\sqrt{2} - 1) \right)^4$$

$$\text{Res}(gf^4)\Big|_{z=z_2} = \frac{1}{4} \left( \frac{3}{2} \pi i + \ln(\sqrt{2} - 1) \right)^4, \quad \text{Res}(gf^4)\Big|_{z=z_3} = \frac{1}{4} \left( \frac{3}{2} \pi i + \ln(\sqrt{2} + 1) \right)^4$$

As above the terms multiplied by the imaginary unit  $i$  in (\*) is equal to zero. By the formula

$$\begin{aligned} \frac{1}{4} \left( \frac{3}{2} \pi i + \ln(\sqrt{2} - 1) \right)^4 &= \frac{1}{4} \left( \frac{81}{16} \pi^4 + \ln^4(\sqrt{2} - 1) - 4 \frac{27}{8} \pi^4 i \ln(\sqrt{2} - 1) + \right. \\ &\left. + 4 \frac{3}{2} \pi i \ln^3(\sqrt{2} - 1) + 6 \frac{-9}{4} \pi^2 \ln^2(\sqrt{2} - 1) \right) \end{aligned}$$

and by summing all the residues, the nonzero terms we get

$$\begin{aligned} &\pi^4 \left( -1 + \frac{81}{64} + \frac{81}{64} + \frac{1}{64} + \frac{1}{64} \right) + \\ &\pi^2 \left( -\frac{27}{8} \ln^2(\sqrt{2} - 1) - \frac{27}{8} \ln^2(\sqrt{2} + 1) - \frac{3}{8} \ln^2(\sqrt{2} - 1) - \frac{3}{8} \ln^2(\sqrt{2} + 1) \right) + \\ &+ \left( \frac{1}{4} \ln^4(\sqrt{2} - 1) + \frac{1}{4} \ln^4(\sqrt{2} - 1) + \frac{1}{4} \ln^4(\sqrt{2} + 1) + \frac{1}{4} \ln^4(\sqrt{2} + 1) \right) = \\ &= \frac{25}{16} \pi^4 - \frac{15}{2} \pi^2 \ln^2(\sqrt{2} + 1) + \ln^4(\sqrt{2} + 1) \end{aligned}$$

The final computation is

$$I = \frac{1}{4} \left( - \sum \text{Res}(g(z)f^4(z)) + 16\pi^2 \int_{-1}^1 gf \, dx \right)$$

namely

$$\frac{1}{4} \left( -\frac{25}{16} \pi^4 + \frac{15}{2} \pi^2 \ln^2(\sqrt{2} + 1) - \ln^4(\sqrt{2} + 1) + 16\pi^2 \left( \frac{\pi^2}{8} - \frac{1}{2} (\ln^2(1 + \sqrt{2})) \right) \right)$$

and this yields

$$I = \frac{7}{64} \pi^4 - \frac{\pi^2}{8} \ln^2(\sqrt{2} + 1) - \frac{1}{4} \ln^4(\sqrt{2} + 1)$$

*Also solved by Yunyong Zhang, Chinaunicom; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Theo Koupelis, Clark College, WA, USA; Seán M. Stewart, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia.*

## Olympiad problems

O625. Find all primes  $p$  such that

$$\frac{(p-2)! - 1}{p^2} = \frac{2(p^4 + 3p^2 - 9)}{p-1}.$$

*Proposed by Titu Andreescu, University of Texas at Dallas, USA*

*Solution by Emon Suin, Ramakrishna Mission Vidyalaya, Narendrapur, West Bengal, India*

We first have from the consequence of the Wilson's Theorem,  $(p-2)! \equiv 1 \pmod{p}$  and hence,  $p$  divides  $(p-2)! - 1$ . Let  $(p-2)! - 1 = pk$ . Then,

$$\begin{aligned} \frac{kp}{p^2} &= \frac{2(p^4 + 3p^2 - 9)}{p-1} \\ \iff k(p-1) &= 2p(p^4 + 3p^2 - 9). \end{aligned}$$

Now, since  $\gcd(p, p-1) = 1$ ,  $k$  must be divisible by  $p$ . So, the left hand side of the given equation is an integer, and hence, the right hand side must be an integer as well. Thus,  $p-1 \mid 2(p^4 + 3p^2 - 9)$ , which can be written as  $2(p^2 + 9)(p^2 - 1) - 10p^2$ . Since  $p-1 \mid (p^2 - 1)$ , we must have  $10p^2$  to be divisible by  $p-1$ . As  $\gcd(p, p-1) = 1$ ,  $p-1 \mid 10$ , and hence,  $p = 11$  is the only solution.

*Also solved by Soham Dutta, West Bengal, India; Batakogias Panagiotis, High School of Velestino, Greece; Anmol Kumar, IISc, Bangalore, India; Ivan Hadinata, Gadjah Mada University, Yogyakarta, Indonesia; Sundaresh H R, Shivamogga, India; Theo Koupelis, Clark College, WA, USA.*

O626. Prove that there are infinitely many triples  $(a, b, k)$  of positive integers such that

$$\frac{a+1}{b} + \frac{b+1}{a} = k$$

and find all possible values of  $k$ .

*Proposed by Mircea Becheanu, Canada*

*Solution by the author*

We start by considering particular cases.

*Case 1:* Assume that  $a = b$ . The equation becomes

$$2\left(\frac{a+1}{a}\right) = k \Leftrightarrow \frac{2}{a} = k - 2$$

and this requires  $a|2$ . We get cases:

*Case (1.i):*  $a = b = 1$  giving the triple  $(a, b, k) = (1, 1, 4)$ .

*Case (1.ii):*  $a = b = 2$  giving the triple  $(a, b, k) = (2, 2, 3)$ .

Therefore we can assume that  $a > b$ . In this situation we consider two cases.

*Case 2:* Assume that  $a = b + 1$ . The equation becomes

$$\frac{b+2}{b} + 1 = k \Leftrightarrow \frac{2}{b} = k - 2,$$

giving that  $b|2$ . We get cases

*Case (2.i):*  $b = 1, a = 2$  which gives the triple  $(a, b, k) = (2, 1, 4)$ .

*Case (2.ii):*  $b = 2, a = 3$  which gives the triple  $(a, b, k) = (3, 2, 3)$ .

*Case 3:* Assume that  $a > b + 1$ . The given equation can be written as a quadratic equation in  $a$ :

$$a^2 - (kb - 1)a + b^2 + b = 0. \tag{1}$$

We apply Fermat descend method. Assume that  $(a, b, k)$  is a triple which satisfies (1) in which  $a > b + 1$ . Such triple exists like, for example  $(14, 6, 3)$  or  $(21, 6, 4)$ , in which both cases  $k = 3$  and  $k = 4$  appear. The quadratic equation (1) has roots  $a_1, a_2$  in which  $a_1 = a$ . Then

$$a_2 = kb - 1 - a = \frac{b(b+1)}{a}$$

is also a root of (1). We prove that  $a_2 < b$ :

$$a_2 < b \Leftrightarrow \frac{b(b+1)}{a} < b \Leftrightarrow b(b+1) < a^2$$

and this is true, because  $a > b + 1 > b$ . So, the triple  $(b, a_2, k)$  is a solution of the problem with  $b > a_2$  and in which  $b + a_2 < a + b$ . If  $b > a_2 + 1$  we can apply again descend method to obtain a new triple  $(a_2, b_1)$  in which  $a_2 > b_1$  and  $a_2 + b_1 < b + a_2$ . So, starting from  $a_1 = a, b_1 = b, k$  we can construct a going down sequence of triples

$$\dots \leftarrow (a_3, b_3, k) \leftarrow (a_2, b_2, k) \leftarrow (a_1, b_1, k) \tag{2}$$

in which  $a_i > b_i$  and the sums  $a_i + b_i$  decrease. The sequence can be continued as long as  $a_i > b_i + 1$ . This can not be done indefinitely, so at some step we arrive to a triple  $(a_i, b_i, k)$  in which  $a_i = b_i + 1$ . But in this case we have one of the cases (2.i) or (2.ii) which show that  $k = 3$  or  $k = 4$ .

The transformations from the sequence (2) can also be applied up, so we obtain infinitely many triples.

*Also solved by Theo Koupelis, Clark College, WA, USA; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania.*

O627. Prove that 3 is the largest value of the positive constant  $k$  such that

$$\frac{1}{ab+k} + \frac{1}{bc+k} + \frac{1}{cd+k} + \frac{1}{da+k} \geq \frac{4}{1+k}$$

for all nonnegative real numbers  $a, b, c, d$  satisfying  $ab + ac + ad + bc + bd + cd = 6$ .

*Proposed by Vasile Cîrtoaje, Oil-Gas University, Ploiești, România*

*Solution by the author*

By choosing  $b = d = \sqrt{ac}$ , the constraint becomes  $(a + c)\sqrt{ac} + ac = 3$ , while the inequality can be written as follows:

$$\frac{1}{a\sqrt{ac}+k} + \frac{1}{c\sqrt{ac}+k} \geq \frac{2}{1+k},$$

$$\frac{(a+c)\sqrt{ac}+2k}{a^2c^2+k(a+c)\sqrt{ac}+k^2} \geq \frac{2}{1+k}.$$

Denoting  $x = ac$ , we have  $3 = (a + c)\sqrt{ac} + ac \geq 3ac = 3x$ , hence  $x \in (0, 1]$ . Since  $(a + c)\sqrt{ac} = 3 - x$ , the inequality becomes

$$\frac{3+2k-x}{x^2-kx+k^2+3k} \geq \frac{2}{1+k},$$

which is equivalent to  $(x-1)(2x+3-k) \leq 0$ . Since  $x-1 \leq 0$ , the inequality is true if and only if  $2x+3-k \geq 0$  for  $x \in (0, 1]$ . So, we get the necessary condition  $3-k \geq 0$ , that is  $k \leq 3$ .

To show that 3 is the largest positive  $k$ , we need to prove the inequality

$$\frac{1}{ab+3} + \frac{1}{bc+3} + \frac{1}{cd+3} + \frac{1}{da+3} \geq 1.$$

By the AM-GM inequality, we have

$$6 = ab + ac + ad + bc + bd + cd \geq 6\sqrt[6]{a^3b^3c^3d^3},$$

hence

$$abcd \leq 1.$$

Write the required inequality as follows:

$$\left(\frac{1}{ab+3} + \frac{1}{cd+3}\right) + \left(\frac{1}{bc+3} + \frac{1}{da+3}\right) \geq 1,$$

$$\frac{3(ab+cd)+18}{abcd+3(ab+cd)+9} + \frac{3(bc+da)+18}{abcd+3(bc+da)+9} \geq 3,$$

$$1 + \frac{9-abcd}{abcd+3(ab+cd)+9} + 1 + \frac{9-abcd}{abcd+3(bc+da)+9} \geq 3,$$

$$\frac{1}{abcd+3(ab+cd)+9} + \frac{1}{abcd+3(bc+da)+9} \geq \frac{1}{9-abcd}.$$

According to the AM-HM inequality, it is sufficient to show that

$$\frac{4}{[abcd+3(ab+cd)+9] + [abcd+3(bc+da)+9]} \geq \frac{1}{9-abcd},$$

which is equivalent to

$$6 \geq ab + bc + cd + da + 2abcd,$$

$$ac + bd \geq 2abcd.$$

Indeed, we have

$$ac + bd \geq 2\sqrt{abcd} \geq 2abcd.$$

The proof is completed. For  $k \in (0, 3]$ , the equality occurs when  $a = b = c = d = 1$ .

*Also solved by Theo Koupelis, Clark College, WA, USA; Emon Suin, Ramakrishna Mission Vidyalaya, Narendrapur, West Bengal, India.*



O628. Let  $x, y, z$  be positive real numbers such that  $x^2 + y^2 + z^2 + xyz = 4$ . Prove that

$$(x + y + z - 2) \left( \frac{1}{x} + \frac{1}{y} + \frac{1}{z} - \frac{1}{2} \right) \geq \frac{5}{2}.$$

*Proposed by Marius Stănean, Zalău, România*

*Solution by the author*

The inequality is equivalent to

$$\frac{(x + y + z - 2) [2(xy + yz + zx) - xyz]}{2xyz} \geq \frac{5}{2},$$

$$(x + y + z - 2)^2(x + y + z + 2) \geq 5xyz.$$

Let  $x = 2 \cos A$ ,  $y = 2 \cos B$ ,  $z = 2 \cos C$ , where  $ABC$  is an acute triangle. The inequality becomes

$$(\sum \cos A - 1)^2 (\sum \cos A + 1) \geq 5 \prod \cos A,$$

inequality, which is obviously true for a triangle that is not acute. Using known identities

$$\cos A + \cos B + \cos C = 1 + \frac{r}{R}, \quad \cos A \cos B \cos C = \frac{s^2 - (2R + r)^2}{4R^2}$$

we get

$$\frac{r^2(2R + r)}{R^3} \geq \frac{5(s^2 - r^2 - 4Rr - 4R^2)}{4R^2},$$

$$\frac{5s^2}{R^2} \leq \frac{4r^3}{R^3} + \frac{13r^2}{R^2} + \frac{20r}{R} + 20.$$

Using the Ravi's substitutions i.e.  $a = y + z$ ,  $b = z + x$ ,  $c = x + y$ ,  $x, y, z > 0$ , the inequality can be rewritten as follows

$$\frac{5 \cdot 16xyz(x + y + z)^3}{(x + y)^2(y + z)^2(z + x)^2} \leq$$

$$\frac{4 \cdot 4^3 x^3 y^3 z^3}{(x + y)^3(y + z)^3(z + x)^3} + \frac{13 \cdot 4^2 x^2 y^2 z^2}{(x + y)^2(y + z)^2(z + x)^2} + \frac{20 \cdot 4xyz}{(x + y)(y + z)(z + x)} + 20,$$

or

$$5P^3 + 20xyzP^2 + 52x^2y^2z^2P + 64x^3y^3z^3 - 20xyz(x + y + z)^3P \geq 0,$$

where we denote  $P = (x + y)(y + z)(z + x)$ . After expanding and reducing terms, it becomes

$$5[(6, 3, 0)] - 5[(6, 2, 1)] + 15[(5, 4, 0)] - 15[(5, 2, 2)] +$$

$$5[(4, 4, 1)] - 13[(4, 3, 2)] + 8[(3, 3, 3)] \geq 0.$$

By Muirhead's Inequality, we have  $[(6, 3, 0)] \geq [(6, 2, 1)]$  and  $[(5, 4, 0)] \geq [(5, 2, 2)]$ , so it remains to show that

$$3[(5, 4, 0)] - 3[(5, 2, 2)] + 5[(4, 4, 1)] - 13[(4, 3, 2)] + 8[(3, 3, 3)] \geq 0. \quad (1)$$

Without loss of generality, we may assume that  $z = \max\{x, y, z\}$  and let denote  $M = (x - y)^2$ ,  $N = (x - z)(y - z) \geq 0$ . We have

$$[(5, 4, 0)] - [(5, 2, 2)] =$$

$$M[z^5(x + y)^2 + z^3(x^4 + x^3y + x^2y^2 + xy^3 + y^4) + z^2xy(x^3 + x^2y + xy^2 + y^3)$$

$$- zx^2y^2(x^2 + xy + y^2) - x^3y^3(x + y)] + N[x^5(y + z)^2 + y^5(z + x)^2].$$

$$\begin{aligned}
& 5[(4, 4, 1)] - 13[(4, 3, 2)] + 8[(3, 3, 3)] = \\
& 10xyz(x^3y^3 + y^3z^3 + z^3x^3 - 3x^2y^2z^2) - 13x^2y^2z^2 \left( \sum_{cyc} x(y^2 + z^2) - 6xyz \right) = \\
& 10xyz(xy + yz + zx)(z^2M + xyN) - 13x^2y^2z^2 [2zM + (x + y)N].
\end{aligned}$$

Returning to (1), the inequality can be rewritten as follows  $\alpha M + \beta N \geq 0$ , where

$$\begin{aligned}
\alpha &= 3z^5(x + y)^2 + 3z^3(x^4 + x^3y + x^2y^2 + xy^3 + y^4) + 3z^2xy(x^3 + x^2y + xy^2 + y^3) \\
&\quad - 3zx^2y^2(x^2 + xy + y^2) - 3x^3y^3(x + y) + 2xyz^3(5yz + 5zx - 8xy) \\
&= 3z^5(x + y)^2 + 3z^2xy(x^3 + x^2y + xy^2 + y^3) + 3(z^3xy - zx^2y^2)(x^2 + xy + y^2) \\
&\quad + 3x^4(z^3 - y^3) + 3y^4(z^3 - x^3) + 2xyz^3(5yz + 5zx - 8xy) \geq 0,
\end{aligned}$$

and

$$\begin{aligned}
\beta &= 3x^5(y + z)^2 + 3y^5(z + x)^2 + x^2y^2z(10xy - 3zx - 3yz) \\
&= 3z^2(x^5 + y^5 - x^3y^2 - x^2y^3) + 2xyz(3x^4 + 5x^2y^2 + 3y^4) + 3x^2y^2(x^3 + y^3) \geq 0,
\end{aligned}$$

which is clearly true.

*Also solved by Paolo Perfetti, Università degli studi di Tor Vergata Roma, Italy; Arkady Alt, San Jose, CA, USA; Nicușor Zlota, Traian Vuia Technical College, Focșani, Romania; Theo Koupelis, Clark College, WA, USA.*

O629. Let  $a, b, c, d$  be positive integers such that  $4(a^2 + b^2) = 5(c^2 + d^2)$  and  $ad - bc$  divides  $c^2 + d^2$ . Prove that

$$2(ac + bd) = L_{2n+1}|ad - bc|$$

for some nonnegative integer  $n$ , where  $L_k$  is the  $k^{\text{th}}$  Lucas number defined by  $L_0 = 2$ ,  $L_1 = 1$  and  $L_{k+1} = L_k + L_{k-1}$ ,  $k = 1, 2, 3, \dots$

*Proposed by Titu Andreescu, University of Texas at Dallas, USA*

*Solution by Theo Koupelis, Clark College, WA, USA*

Let  $c^2 + d^2 = (ad - bc)m$ , where  $m$  is an integer  $\neq 0$ . Using the identity  $(ad - bc)^2 + (ac + bd)^2 = (a^2 + b^2)(c^2 + d^2)$  we get

$$\frac{(c^2 + d^2)^2}{m^2} + (ac + bd)^2 = \frac{5}{4}(c^2 + d^2)^2,$$

and thus

$$4m^2(ac + bd)^2 = (c^2 + d^2)^2(5m^2 - 4) = (ad - bc)^2 m^2(5m^2 - 4),$$

or

$$2(ac + bd) = |ad - bc| \cdot \sqrt{5m^2 - 4}.$$

It is well known, however, that the solutions of the Diophantine equation  $5m^2 - 4 = s^2$ , with  $s$  a positive integer, are given by  $m = \pm F_{2k+1}$  and  $s = L_{2k+1}$ , where  $L_k$  is the  $k^{\text{th}}$  Lucas number, and  $F_k$  is the  $k^{\text{th}}$  Fibonacci number, with  $L_0 = 2$ ,  $L_1 = 1$ ,  $L_{k+1} = L_k + L_{k-1}$ , and  $F_0 = 0$ ,  $F_1 = 1$ , and  $F_{k+1} = F_k + F_{k-1}$ ,  $k = 1, 2, 3, \dots$ . Thus,

$$2(ac + bd) = L_{2n+1}|ad - bc|$$

for some nonnegative integer  $n$ .

O630. Prove that there are infinitely many positive integers  $m$  such that  $m+1, 2m+1, 3m+1$  are all composite and divide  $2^m - 1$ .

*Proposed by Navid Safaei, Sharif University of Technology, Tehran, Iran*

*Solution by the author*

Taking  $m = 6n$ , for some positive integer  $n$ , we are going to prove that there are infinitely many  $n$  such that the three composite numbers  $6n+1, 12n+1, 18n+1$  are composite and divide  $2^{6n} - 1$ .

Let  $n = \frac{2^{36}-1}{9}$  then  $6n+1 = \frac{2^{37}+1}{3}, 12n+1 = \frac{2^{38}-1}{3}, 18n+1 = 2^{37} - 1$  are composite and divide  $2^{6n} - 1$ . Let  $N = \frac{2^{12n}-1}{9}$  then  $6N = \frac{2(2^{12n}-1)}{3}$  thus  $2(6n+1)(12n+1) \mid 6N$ . Hence,

$$\begin{aligned} 6N+1 &= \frac{2^{12n+1}+1}{3} \mid 2^{2(12n+1)} - 1 \mid 2^{6N} - 1, \\ 12N+1 &= \frac{2^{2(6n+1)}-1}{3} \mid 2^{2(6n+1)} - 1 \mid 2^{6N} - 1, \\ 18N+1 &= 2^{12n} - 1 \mid 2^{6N} - 1. \end{aligned}$$

Since  $6N+1, 12N+1, 18N+1$  are all composite and  $N = \frac{2^{12n}-1}{9} > n$  we are done.

Notice that

$$\begin{aligned} 6n+1 \mid 2^{6n} - 1 \mid 2^{6n+1} - 2, \\ 12n+1 \mid 2^{6n} - 1 \mid 2^{12n+1} - 2, \\ 18n+1 \mid 2^{6n} - 1 \mid 2^{18n} - 1 \mid 2^{18n+1} - 2. \end{aligned}$$

*Remark:* Let by  $t_n$  and  $\omega_n$  we denote the  $n$ -th triangular and pentagonal numbers, we can also prove  $t_{12n+1} = (6n+1)(12n+1) \mid 2^{6n} - 1 \mid 2^{t_{12n+1}} - 2$ . Let  $\omega_{12n+1} = (12n+1)(18n+1)$  then  $\omega_{12n+1} \mid 2^{\omega_{12n+1}} - 2$ .

*Also solved by Zijin Peng, The Affiliated High School of South China Normal University, Guangzhou, Guangdong, China.*